

**XV REUNIÓN CIENTÍFICA
SOCIEDAD ESPAÑOLA DE CROMATOGRFÍA
Y TÉCNICAS AFINES · SECyTA 2015**

*XV SCIENTIFIC MEETING
OF THE SPANISH SOCIETY OF CHROMATOGRAPHY
AND RELATED TECHNIQUES · SECyTA2015*

**VII REUNIÓN NACIONAL
SOCIEDAD ESPAÑOLA DE ESPECTROMETRÍA DE MASAS · SEEM 2015**

*VII NATIONAL MEETING
OF THE SPANISH SOCIETY OF MASS SPECTROMETRY · SEEM2015*

CASTELLÓN
UNIVERSITAT JAUME I

27-30 de octubre de 2015

October 27-30, 2015

www.secytaseem2015.com



Organiza
Organizer

SECyTA

SOCIEDAD ESPAÑOLA DE
CROMATOGRFÍA
Y TÉCNICAS AFINES

S.E.E.M. 

Colabora
Collaborate



**UNIVERSITAT
JAUME I**
DES DE 1991

25
ANYS

COMITÉS

COMITÉ CIENTÍFICO

Esteban Abad Holgado, Institut de Diagnòstic Ambiental i Estudis de l'Aigua (CSIC)
Joaquín Beltrán Arandes, Universitat Jaume I de Castelló
Jordi Díaz Ferrero, Instituto Químico de Sarrià, Universitat Ramón Llull
Ana M^a García Campaña, Universidad de Granada
María José González Carlos, Instituto de Química Orgánica General (CSIC)
Belén Gómara Moreno, Instituto de Química Orgánica General (CSIC)
Joan Grimalt Obrador, Institut de Diagnòstic Ambiental i Estudis de l'Aigua (CSIC)
Félix Hernández Hernández, Universitat Jaume I de Castelló
Elena Ibáñez Ezequiel, Instituto de Investigación en Ciencias de la Alimentación (CSIC)
Begoña Jiménez Luque, Instituto de Química Orgánica General (CSIC)
Encarnación Moyano Morcillo, Universitat de Barcelona
Yolanda Picó Garcia, Universitat de València
Juan Vicente Sancho Llopis, Universitat Jaume I de Castelló
Fco. Javier Santos Vicente, Universitat de Barcelona
M^a Luz Sanz Murías, Instituto de Química Orgánica General (CSIC)
Rosa Ventura Alemany, Institut Hospital del Mar d'Investigacions Mèdiques (IMIM)

COMITÉ ORGANIZADOR

Juan Vicente Sancho Llopis, Universitat Jaume I de Castelló (Chair)
Joaquín Beltrán Arandes, Universitat Jaume I de Castelló (Co-Chair)
Esteban Abad Holgado, Institut de Diagnòstic Ambiental i Estudis de l'Aigua (CSIC)
Jordi Díaz Ferrero, Instituto Químico de Sarrià, Universitat Ramón Llull
María José González Carlos, Instituto de Química Orgánica General (CSIC)
Belén Gómara Moreno, Instituto de Química Orgánica General (CSIC)
Maria Ibáñez Martínez, Universitat Jaume I de Castelló
Francisco López Benet, Universitat Jaume I de Castelló
Encarnación Moyano Morcillo, Universitat de Barcelona
Elena Pitarch Arquimbau, Universitat Jaume I de Castelló
Tània Portolés Nicolau, Universitat Jaume I de Castelló
Antoni Francesc Roig i Navarro, Universitat Jaume I de Castelló
Roque Serrano Gallego, Universitat Jaume I de Castelló
Juan Solé Ribalta, HiTIC, S.A.
Rosa Ventura Alemany, Institut Hospital del Mar d'Investigacions Mèdiques (IMIM)

P-49

LEVELS AND TEMPORAL TRENDS OF PHTHALATE METABOLITES IN HUMAN SPANISH URINE SAMPLES FROM 2012 TO 2015

Mario A. Fernández^{(1)*}, Laura Herrero⁽¹⁾, Jesús E. Quintanilla⁽²⁾, Sagrario Calvarro⁽¹⁾, Belén Gómara⁽¹⁾, María José González⁽¹⁾.

⁽¹⁾ General Organic Chemistry Institute (CSIC). Juan de la Cierva 3, 28006-Madrid, Spain.

⁽²⁾ Institute of Physical Chemistry "Rocasolano" (CSIC). Serrano 119, 28006-Madrid, Spain

*mario@iqog.csic.es, Tel: +34 91 561 88 06, Fax: +34 564 48 53

Phthalates (diesters of 1,2-benzenedicarboxylic acid), also known as phthalic acid esters (PAEs), are synthetic organic compounds that are used in a broad spectrum of industrial and commercial applications, like toys, food packaging, lubricants, vinyl flooring, adhesives, detergents, hair spray, shampoo, etc. ^[1]. They have endocrine disrupting potency and have become nearly ubiquitous. For the general population, the human exposure to phthalates is through ingestion, inhalation, dermal absorption, and/or contact with medical devices ^[2]. Once they are ingested, PAEs are rapidly metabolized to their respective monoesters in human body and some of these primary metabolites can be further metabolized to secondary metabolites and excreted by urine. All of these metabolites can be used as biomarkers of human exposure to phthalates ^[3].

The sample preparation and instrumental determination methods have been previously described elsewhere ^[4]. Briefly, the sample preparation consisted of an enzymatic deconjugation of the glucuronide phthalates followed by a purification step with OASIS[®] HLB SPE cartridge (Waters, MA, USA). The instrumental determination was carried out by using UPLC QqQ MS² (Xevo-TQS, Waters) working in MRM mode, with a BEH-phenyl column (50 mm x 2.1 mm, 1.7 µm),.

We present here, for the first time, the levels and temporal trends of 9 PAE metabolites (6 phthalate monoesters (monomethyl phthalate [MMP], monoethyl phthalate [MEP], mono-*iso*-buthyl phthalate [MiBP], mono-*n*-butyl phthalate [MBP], monobenzyl phthalate [MBzP] and monoethylhexyl phthalate [MEHP]) and 3 secondary metabolites of DEHP (5-OH-mono(2-ethylhexyl) phthalate [5-OH-MEHP], 5-oxo-mono(2-ethylhexyl) phthalate [5oxo-MEHP] and 5-carboxy-mono(2-ethylpentyl) phthalate [5cx-MEPP]) in human urine samples collected in 2012 and 2015 from 9 volunteers living in the Community of Madrid. Detectable amounts of the 9 MPAEs were obtained in all the studied samples, being the MPAE levels found in almost all collected in 2012 slightly lower than those found three years later (2015).

[1] T. Schettler. *Int. J. Androl.* **29** (2006) 134-139.

[2] R. Hauser, A. Calafat. *Occup. Environ. Med.* **62** (2005) 806-818.

[3] H. Frederiksen, S.K. Kranich. *Environ. Sci. Technol.* **47** (2013) 958-967.

[4] L. Herrero, S. Calvarro. *Anal Chim Acta* **853** (2015) 625-636.

Acknowledgements: Authors wish to thank all the volunteers who have donated samples and Paula Chamorro and Héctor Carrascosa for their laboratory work. Financial support was obtained from the Spanish Ministry of Economy and Competitiveness (project AGL2012-37201) and the Comunidad Autónoma of Madrid and European funding from FEDER program (projects S2009/AGR-1464, ANALISYC-II and S2013/ABI-3028, AVANSECAL).